

منابع:

۱. شجاع، محمود و هوشنگ ساعدی. (۱۳۷۵). گیاهان سمی و تاثیر مسمومیت آنها در حیوانات. انتشارات دانشگاه تهران.
۲. مشیدی، بدیع الله. و مهدی سمسار. (۱۳۵۲). مفردات پزشکی. انتشارات دانشگاه تهران.
3. Ambros, P.F. , Matzke, A.J.M. and M.A. Matzke. (1986). Localization of *Agrobacterium rhizogenes* T-DNA in plant chromosomes by *in situ* hybridization. *EMBO J.* 19: 421-432.
4. Archana, G. and M.L. Narasu. (2000). Transgenic hairy roots: Recent trends applications: *a review. Biotechnology Advances.* 18: 1-22.
5. Balandrin, M.F. , Klocke, J.A. , Wurtele, E.S. and W.H. Bollinger. (1985). Natural plant chemicals: sources of industrial and medicinal materials. *Science* 228: 1154-1160.
6. Bensaddek, L. , Gillet, F. , Saucedo, J.E.N. and M.A. Fliniaux. (2001). The effect of nitrate and ammonium concentration on growth and alkaloid accumulation of *Atropa belladonna* hairy roots. *J. Biotech.* 85: 35-40
7. Blackshaw, R.E. (1998). Weed management in beans. Agriculture and AgrifoodCanada, Lethridge. WWW.Pulse.A/B.Ca/Newsletter/98 spring/bean.html.
8. Bonhomme, V. , Laurain-Mattar, D. , Lacoux, J. , Fliniaux, M.A. and A. Jacquin-Dubreuil. (2000). Tropan alkaloid production by hairy roots of *Atropa belladonna* obtained after transformation with *Agrobacterium rhizogenes* 15834 and *Agrobacterium tumefaciens* containing *rol* A, B, C genes only. *J. Biotech.* 81: 151-158.
9. Bourgaud, F. , Gravot, A. , Milesi, S. and E. Gontier. (2001). Production of plant secondary metabolites: a historical perspective. *A Review. Plant Sci.* 161: 839-851.
10. Byrne, M.C. , Koplow, J. , David, C. , Tempe, J. and M.D. Chilton. (1983). Structure of T-DNA in root transformed by *Agrobacterium rhizogenes*. *J. Appl. Genet.* 2:201-209.
11. Carderelli, M. , Mariotti, D. , Pomponi, M. , Spano, L. , Capone, I and P. Castantino. (1987). *Agrobacterium rhizogenes* T-DNA genes capable of inducing hairy root phenotype. *Mol. Genet.* 209: 475-480.
12. Collins, D.J. , Culvenor, C.C.J. , Lamberton, J.A. , Loder, J.W. and J.R. Price. (1990). Plants for Medicines. CSIRO, Melbourne.
13. Cusido, R.M. , Palazon, J. , Pinol, M.T. , Bonfill, M. and C. Morales. (1998). *Datura metel*: In vitro production of tropan alkaloids. *Planta Medica.* 65: 144-148.
14. DellaPenna, D. (2001). Plant metabolic engineering. *Plant Physiol.* 125: 160-163.
15. Dixon, R.A. (1999). Flavonoids and isoflavonoids a goldmine for metabolic engineering. *Trend Plant Sci.* 4: 394-400.
16. Dixon, R.A. (2001). Natural products and plant disease resistance. *Nature.* 411: 843-847.
17. Doran, P.M. (1989). Prospects for production of plant chemicals from genetically transformed roots. *Aust. J. Biotechnol.* 3: 270-277.

18. Erickson, D. (1991). Secret garden. *Sci. Amer.* **265**: 121-122.
19. Facchini, P.J. (2001). Alkaloid biosynthesis in plant: biochemistry, cell biology, molecular regulation, and metabolic engineering applications. *Annu Rev Plant Physiol.* **52**: 29-66.
20. Flores, H.E. , Hoy, M.W. and J.J. Pickard. (1987). Secondary metabolites from root culture. *J. Biotechnol.* **5**: 64-69.
21. Forkmann, G. and S. Martens. (2001). Metabolic engineering and application of flavonoids. *Curr Opin Biotechnol.* **12**: 155-160.
22. Frick, B. and A.G. Thomas. (1992). Weed surveys in different tillage, systems in southwestern ontario field. *Crops.Can.Journal. Plant Science.* **72**:1337-1347.
23. Fujita, Y. (1990). The production of industrial compounds. In: Bhojwani, S.S. (ed.). *Plant Tissue Culture: Applications and Limitations.* Elsevier, Amsterdam (pp: 259-275).
24. Giri, A. , Banerjee, S. , Ahuja, P.S. and C.C. Giri. (1997). Production of hairy roots in *Aconitum heterophyllum* wall using *Agrobacterium rhizogenes*. In *Vitro Cell Dev Biol-Plant.* **33**: 280-283.
25. Hamill, J.D. , Parr, A.J. , Rhodes, M.J.C. , Robins, R.J. and N.J. Walton (1987). New routes to plant secondary products. *Biotechnol.* **5**: 800-804.
26. Hamill, J.D. (1993). Alterations in auxin and cytokinin metabolism of higher plants due to expression of specific genes from pathogenic bacteria: a review. *Aust. J. Plant Physiol.* **20**: 405-423.
27. Hamill, J.D. and M.C.J. Rhodes. (1993). Manipulation secondary metabolism in culture. In: Grierson, D. (ed.). *Plant Biotechnology.* Blackie Academic and Professional, London. Vol 3 (pp: 178-209).
28. Hamill, J.D. , Parr, A.J. , Robins, R.J. and M.C.J. Rhodes (1986). Secondary product formation by cultures of *Beta vulgaris* and *Nicotiana rustica* transformed with *Agrobacterium rhizogenes*. *Plant Cell Rep.* **5**: 111-114.
29. Harborne, J.B. (1999). Classes and functions of secondary products. In: Walton, N.J. and D.E. Brown, (ed.). *Chemicals from Plants, Perspectives on Secondary Plant Products.* Imperial College Press. (pp: 1-25).
30. Hashimoto, T. , Yun, D.J. and Y. Yamada. (1993). Production of tropane alkaloids in genetically engineered root cultures. *Phytochemistry.* **32**: 713-718.
31. Heble, M.R. (1985). Multiple shoot cultures: a viable alternative *in vitro* system for the production of known and new biologically active plant constituents. In: Neumann, K.H. Barz, W. and E. Reinhard, (eds). *Primary and Secondary Metabolism of Plant Cell Cultures.* Springer-Verlag, Berlin (pp: 281-289).
32. Hein, M.B. , Tang, Y. , McLeod, D.A. , Janda, K.D. and A. Hiatt (1991). Evaluation of immunoglobulins from plant cells. *Biotechnol. Prog.* **7**: 455-461.
33. Hilton, M.G. and M.J. Rhodes. (1990). Growth and hyoscyamine production of hairy root cultures of *Datura stramonium* in a modified stirred tank reactor. *Appl Microbiol Biotechnol.* **33**: 132-138.
34. Hiraoka, N and M, Tabata. (1974). Alkaloid production by plants regenerated from cultured cells of *Datura innoxia*. *Phytochemistry.* **13**:1671-1675.
35. Irena, R. and J. Swanton. (2001). Understanding Maize-weed competition: resource competition, light quality and the whole plant. *Field Crops Research.* **71**:139-150.

36. Jouanin, L. , Guerche, P. , Pamboukgjian, N. , Tourneur, C. , Casse Delbart, F. and J. Tourneur. (1987). Structure of T-DNA in plants regenerated from roots transformed by *Agrobacterium rhizogenes* A4. *Mol. Genet.* **206**: 387-392.
37. Jouhikainen, L.L. , Jokelainen, T. , Hiltunen, R. , Teeri, T.H. and K.M. Oksman-Caldentey. (1999). Enhancement of scopolamine production in *Hyoscyamus muticus* L. hairy root cultures by genetic engineering. *Planta.* **208**: 545-551.
38. Jung, G. and D, Tepfer. (1987). Use of genetic transformation by the Ri T-DNA of *Agrobacterium rhizogenes* to stimulate biomass and tropane alkaloid production in *Atropa belladonna* and *Calystegi sepium* roots grown in vitro. *Plant Sci.* **50**:145-151.
39. Korber, H. , Strizhov, N. , Staiger, D. , Feldwisch, J. , Olsson, O. , Sandberg, G. , Palme, K. , Schell, J. and C. Koncz. (1991). T-DNA gene 5 of *Agrobacterium* modulates auxin response by autoregulated synthesis of a growth hormone antagonist in plant. *EMBO J.* **10**: 3983-3991.
40. Krolicka, A. , Stabisewska, I. , Bielaeski, K. , Malinski, E. , Szafranek, J. and E, Ljkowska. (2001). Establishment of hairy root cultures of *Ammi majus*. *Plant Sci.* **160**:259-264.
41. Kutchan, T.M. (1995). Alkaloid biosynthesis the basis for metabolic engineering of medicinal plants. *Plant Cell.* **7**: 1059-1070.
42. Li, B.H. , Zhang, H.M. , Xu, T.F. and R.X. Ding. (2000). Genetic transformation of autotetraploid *Isatis indigotica* induced by Ri T-DNA and plant regeneration. *Zhongguo Zhong Yao Za Zhi.* **25**: 657-660
43. Li, J. , Ou-Lee, T.M. , Raba, R. , Amundson, R.G. and R.L. Last. (1993). *Arabidopsis* mutants are hypersensitive to UV-B radiation. *Plant Cell.* **5**: 171-179.
44. Matsubara, K. and Y. Fujita. (1991). Production of berberine. In: Komamine, A. Misawa, M. and F. DiCosmo, (eds). *Plant Cell Culture in Japan*. Ltd. Tokyo. Japan. (pp: 39-44).
45. Menze, A. and C. Ilers. (1999). Transformation of different *Brassica napus* cultivars with three different strains of *Agrobacterium rhizogenes*. *International Rapeseed congress. Australia.*
46. Moyano, E. , Fornale, S. , Palazon, J. , Cusido, R.M. , Bagni, N. and M.T. Pinol. (2002). Alkaolid production in *Duboisia* hybrid hairy root cultures overexpressing the *pmt* gene. *Phytochemistry.* **59**: 697-702.
47. Moyano, E. , Fornale, S. , Palazon, J. , Cusido, R.M. , Bonfil, M. , Morales, C. and M.T. Pifiol. (1999). Effect of *Agrobacterium rhizogenes* T-DNA on alkaloid production in solanaceae plants. *Phytochemistry.* **52**:1287-1299.
48. Muir, S.R. , Collins, G.J. , Robinson, S. , Hughes, S. , Bovy, A. , DeVos, C.H.R. , Van Tunen, A.J. and M.E. Verhoeyen. (2001). Over expression Of *Petunia* chalcone isomerase in tomato results in fruits containing increased levels of flavonols. *Nat Biotechnol.* **19**: 470-474.
49. Mukundan, U. , Rai, A. , Dawda, H. , Ratnaparkhi, S. and V. Bhide. (1998). Secondary metabolites in *Agrobacterium rhizogenes* mediated transformed root cultures. In: Srivastava, P.S. (ed). *Plant Tissue Culture and*

- Molecular Biology Applications and Prospects. New Delhi, Narosa Publishing House. (pp. 302-333).
50. Nussbbaumer, P. , Kapentanidis, I. and P. Christen. (1998). Hairy root of *Datura candida* × *D. Aurea*: Effect of culture medium composition on growth and alkaloid biosynthesis. *Plant Cell Rep.* 17: 405-413.
 51. Palazon, J. , Altabella, T. , Cusido, R.M , Ribo, M. and M.T. Pinol. (1995). Growth and tropane alkaloid production in *Agrobacterium* transformed roots and derived callus of *Datura*. *Biologia Plantarum.* 37: 161-168.
 52. Petiard, V. , Baubault, C. , Baribaud, A. , Hutin, M. and D. Courtois, (1985). Studies on variability of plant tissue cultures for alkaloid production in *Catharanthus roseus* and *Papaver somniferum* callus cultures. In: Neuman, K.H. Barz, W. and E. Reinhard, (eds.). Primary and Secondary Metabolism of Plant Cell Culture. Springer-Verlag, Berlin, Heidelberg, New York. (pp.133-142).
 53. Pinol, M.T. , Palazon, J. , Altabella, T. , Cusido, R. and M. Serrano. (1985). Effect of auxin on alkaloids, K⁺ and free amino acid content in cultured tobacco callus. *Physiol. Plant.* 65:299-304.
 54. Pinol, M.T. , Palazon, J. , Cusido, R.M. and M. Serrano. (1996). Effect of Ri T-DNA from *Agrobacterium rhizogenes* on Growth and hyoscyamine production in *Datura stramonium* root culture. *Botanica.Acta.* 109: 133-138.
 55. Pinol, M.T. , Palazon, J. and M. Serrano. (1984). Growth and nicotine content of tobacco callus cultures without organogenesis. *Plant Sci. Lett.* 36: 219-226.
 56. Ramawat, K.G. and K.M. Merillon. (1999). Biotechnology Secondary Metabolites. Sience Publishers U.S.A.
 57. Rhodes, M.J.C. , Parr, A.J. , Ceielutt, A. and E.L.H. Aird, (1994). Influence of exogenous hormone on the growth and secondary metabolite formation in transformed root cultures. *Plant Cell Tissue and Organ Culture.* 38: 143-193.
 58. Saito, F. , Hashimoto, T. , Hacia, A. , Tamura, K. , Choi, K.B. , Morishinge, T. , Fujimoto, H. and Y. Yamada. (2001). Metabolic engineering of plant alkaloid biosynthesis. *Proc. Nat. Acad. Sci. USA.* 98: 367-372.
 59. Saito, K. , Murakoshi, I. , Inze, D. and M. van Montagu. (1989). Biotransformation of nicotine alkaloids by tobacco shooty teratomas induced by Ti plasmid mutant. *Plant Cell Rep.* 7: 607-610.
 60. Saito, K. , Yamazaki, M. , Kawaguchi, A. and I. Murakoshi. (1991). Metabolism of solanaceous alkaloids in transgenic plant teratomas integrated with genetically engineered genes. *Tetrahedron* 47: 5955-5968.
 61. Sevon, N. , Drager, B. , Hiltunen, R. and K.M. Oksman-Caldentey, (1997). Characterization of transgenic plants derived from hairy roots of *Hyoscyamus muticus*. *Plant Cell Rep.* 16: 605-616.
 62. Sikuli, N.N. and K. Demeyer. (1997). Influence of the ion-composition of the medium on alkaloid production by hairy roots of *Datura stramonium*. *Plant Cell, Tissue and Organ Culture.* 47: 261-267
 63. Spena, A. , Estruch, J.J. , prinsen, E. , Nacken, W. , van onckelen, H. and H. sommer. (1992). Anther specific expression of the rol B gene of *Agrobacterium rhizogenes* increases IAA content in anthers and alters anther development and whole flower growth. *Theor Appl. Genet.* 84:520-527.

64. **Spencer, A. , Hamil, J.D. and M.C.J. Rhodes.** (1990). Production of terpenes by differentiated shoot culture of *Mentha citrata* transformed with *Agrobacterium tumefaciens* T37. *Plant Cell Rep.* **8**: 601-604.
65. **Spencer, A. , Hamil, J.D. and M.C.J. Rhodes.** (1993). In vitro biosynthesis of monoterpenes by *Agrobacterium* transformed shoot cultures of two *Mentha* species. *Phytochem.* **32**: 911-919.
66. **Srinivasan, V. , Pestchanker, L. , Moser, S. , Hirasuna, T.J. , Taticek, R.A. and M.L. Shuler.** (1995). Taxol production in bioreactors: Kinetics of biomass accumulation, nutrient uptake and taxol production by cell suspension of *Taxus baccata*. *Biotechnol Bioeng.* **47**: 666-741.
67. **Taji, A. and W. Richard.** (1996). Tissue Culture of Australian Plants. (pp: 240-283). University of New England, Australia.
68. **Tepfer, D.** (1990). Genetic transformation using *Agrobacterium rhizogenes*. *Physiol. Plant.* **79**: 140-146.
69. **Toivonen, L.** (1993). Utilization of hairy root culture for production of secondary metabolites. *Biotechnol. Prog.* **9**:12-20.
70. **Torssell, K.B.G.** (1997). Chemical ecology. In: Torssell, K.B.G. (ed.). Natural Product Chemistry, A Mechanistic Biosynthetic and Ecological Approach. Swedish Press. (pp: 42-79).
71. **Verpoorte, R. , Vander Heijden, R. and J. Memelink.** (2000). Engineering of the plant cell factory for secondary metabolite production. *Transgenic Rep.* **9**: 323-343.
72. **Verpoorte, R. and J. Memelink.** (2002). Engineering secondary metabolite production in plants. *Biotechnology.* **13**: 181-187
73. **Waller, G.R. and E.k. Nowacki.** (1977). Alkaloid Biology and Metabolism in Plants. Plenum Press. New York.
74. **White, P.R.** (1934). Potentially unlimited growth of excised tomato root tips in a liquid medium. *Plant Physiol.* **9**: 585-600.