

- [1] H. O. Sonkurt, A. E. Altınöz, E. Çimen, F. Köşger, and G. Öztürk, “The role of cognitive functions in the diagnosis of bipolar disorder: A machine learning model,” *Int. J. Med. Inform.*, vol. 145, p. 104311, 2021, doi: 10.1016/j.ijmedinf.2020.104311.
- [2] L. Sierra, Pilar and Livianos, Lorenzo and Arques, Sergio and Castello, Javier and Rojo, “Prodromal symptoms to relapse in bipolar disorder,” *Aust. N. Z. J. Psychiatry*, vol. 41, no. 5, pp. 385–391, 2007, doi: 10.1080/00048670701266854.
- [3] F. Edition, “Diagnostic and Statistical Manual of Mental Disorders (DSM).,” *Am Psychiatr. Assoc.*, vol. 21, 2013.
- [4] Y. Han, Z. Lu, Z. Du, Q. Luo, and S. Chen, “A YinYang bipolar fuzzy cognitive TOPSIS method to bipolar disorder diagnosis,” *Comput. Methods Programs Biomed.*, vol. 158, pp. 1–10, 2018, doi: 10.1016/j.cmpb.2018.02.004.
- [5] L. Singh, L. Bode, C. Budak, K. Kawintiranon, C. Padden, and E. Vraga, “Understanding high- and low-quality URL Sharing on COVID-19 Twitter streams,” *J. Comput. Soc. Sci.*, vol. 3, no. 2, pp. 343–366, 2020, doi: 10.1007/s42001-020-00093-6.
- [6] N. Berry, F. Lobban, M. Belousov, R. Emsley, G. Nenadic, and S. Bucci, “{\#}WhyWeTweetMH: Understanding Why People Use Twitter to Discuss Mental Health Problems,” vol. 19, no. 4, p. e107, Apr. 2017, doi: 10.2196/jmir.6173.
- [7] I. B. Kursuncu, Ugur and Gaur, Manas and Lokala, Usha and Thirunarayan, Krishnaprasad and Sheth, Amit and Arpinar, *Predictive Analysis on Twitter: Techniques and Applications*. Springer, 2019.
- [8] M. hancellor, Stevie and De Choudhury, “Methods in predictive techniques for mental health status on social media: a critical review,” *npj Digit. Med.*, vol. 3, no. 1, pp. 1–11, 2020, doi: 10.1038/s41746-020-0233-7.
- [9] C. Coppersmith, Glen and Dredze, Mark and Harman, *Quantifying Mental Health Signals in Twitter*. Association for Computational Linguistics, 2014.
- [10] G. and Coppersmith, M. and Dredze, C. and Harman, and K. Hollingshead, *From ADHD to SAD: Analyzing the Language of Mental Health on Twitter through Self-Reported Diagnoses*. Association for Computational Linguistics, 2015.
- [11] Y.-S. Saravia, Elvis and Chang, Chun-Hao and De Lorenzo, Renaud Jollet and Chen,

- MIDAS: Mental illness detection and analysis via social media*. IEEE, 2016.
- [12] V. Jarang, Satish and Joshi, Deepali and Deshpande, *Behaviour Analysis Using Word Embedding & Machine Learning on Social Media*. IEEE, 2019.
- [13] J. Beheshti, Amin and Moraveji-Hashemi, Vahid and Yakhchi, Shahpar and Motahari-Nezhad, Hamid Reza and Ghafari, Seyed Mohssen and Yang, *personality2vec : Enabling the Analysis of Behavioral Disorders in Social Networks*. 2020.
- [14] D. Joshi and D. M. Patwardhan, “An analysis of mental health of social media users using unsupervised approach,” *Comput. Hum. Behav. Reports*, vol. 2, no. May, p. 100036, 2020, doi: 10.1016/j.chbr.2020.100036.
- [15] M. Cernak, Š. Beňuš, and A. Lazaridis, “Speech vocoding for laboratory phonology,” *Comput. Speech Lang.*, vol. 42, pp. 100–121, 2017, doi: 10.1016/j.csl.2016.10.001.
- [16] C.-H. Yu, Dong and Siniscalchi, Sabato Marco and Deng, Li and Lee, *Boosting attribute and phone estimation accuracies with deep neural networks for detection-based speech recognition*. IEEE, 2012.
- [17] R. G. Sands, “Bipolar disorder and social work practice.,” *Soc. Work Health Care*, vol. 10, no. 3, pp. 91–105, 1985.
- [18] K. N. Fountoulakis, “The emerging modern face of mood disorders: A didactic editorial with a detailed presentation of data and definitions,” *Annals of General Psychiatry*, vol. 9. 2010, doi: 10.1186/1744-859X-9-14.
- [19] N. Colneric and J. Demsar, “Emotion Recognition on Twitter: Comparative Study and Training a Unison Model,” *IEEE Trans. Affect. Comput.*, vol. 11, no. 3, pp. 433–446, 2020, doi: 10.1109/TAFFC.2018.2807817.